

## LIST OF FIGURES

### Page

Figure 2.1	Location of commercial herring fisheries.....	2- 3
Figure 3.1	Geographical distribution of herring divided into five main groups.....	3- 2
Figure 3.2	Pacific herring.....	3- 3
Figure 3.3	Distribution of Pacific herring. Size of circle approximates range and abundance.....	3- 4
Figure 3.4	Known Pacific herring spawning areas in California.....	3- 7
Figure 3.5	Timing of Pacific herring spawnings by major stock groups.....	3- 8
Figure 3.6	Size distribution of Pacific herring by school in San Francisco Bay during the 1989-90 spawning season.....	3-17
Figure 3.7	Herring growth curves for Tomales and San Francisco Bays.....	3-19
Figure 3.8	Puget Sound food web involving herring. Solid lines indicate primary linkages and dashed lines indicate secondary linkages.....	3-22
Figure 3.9	Occurrence of Pacific herring in the diet of King salmon in the vicinity of San Francisco Bay.....	3-24
Figure 3.10	San Francisco and Tomales Bay spawning stock biomass estimates based on CDFG spawn escape- ment surveys.....	3-33
Figure 3.11	San Francisco spawning stock biomass estimates based on spawn escapement and hydroacoustic surveys.....	3-36
Figure 3.12	San Francisco Bay spawning stock biomass estimates (for 3-yr-old and older) based on	

spawn escapement survey and cohort analysis....3-38

Figure 3.13 Annual landings of Pacific herring in  
California by area.....3-47

Figure 3.14 Commercial fishing districts open to herring  
roe fisheries within San Francisco Bay.....3-55

#### Page

Figure 3.15 Number of permits by gear type issued to fish  
in the San Francisco Bay herring roe fishery...3-56

Figure 3.16 Relationship between targeted catch (20% or  
less of spawning biomass) and actual catch in  
the San Francisco Bay herring roe fishery  
(quota based on previous season biomass).....3-57

Figure 3.17 Areas open to commercial herring roe fishing in  
Tomales and Bodega Bay waters (portion of  
District 10).....3-60

Figure 3.18 Number of permits by gear type issued to fish  
in the Tomales and Bodega Bay herring roe  
fishery.....3-61

Figure 3.19 Relationship between targeted catch (20% or  
less of spawning biomass) and actual catch in  
the Tomales and Bodega Bay herring roe fishery  
(quota based on previous season biomass).....3-62

Figure 3.20 Areas (districts) open to commercial herring  
fishing in ocean waters.....3-66

Figure 3.21 Areas within San Francisco Bay estuary used by  
commercial herring fisheries. Areas marked by  
diagonals conditionally closed to round haul...3-73

Figure 3.22 Areas open to commercial herring fishing in  
Tomales and Bodega Bay waters.....3-85

Figure 3.23 Area open to commercial herring fishing in  
Humboldt Bay waters.....3-88

Figure 3.24 Areas open to commercial herring roe fishing

in the Crescent City area.....	3-93
--------------------------------	------

## LIST OF TABLES

### Page

Table 3.1 Birds Observed Eating Pacific Herring Eggs.....	3-20
Table 3.2 Estimated Number of Younger Herring (x1000) by Year Class in the San Francisco Bay Population..	3-42
Table 3.3 Age Composition of the Tomales Bay Gill Net Catch.....	3-44
Table 4.1 Preliminary Assessment of Environmental Sensitivity to Commercial Herring Fisheries in Selected Areas.....	4- 4
Table 4.2 Daily Emission Rates From Commercial Herring Fishing Vessels (Tons/Day) in Comparison With San Francisco County Emission Rates (1987).....	4- 11
Table 4.3 Daily Emission Rates From Commercial Herring Fishing Support Vehicles (Tons/Day) in Comparison With San Francisco County Emission Rates (1987).....	4- 11
Table 4.4 Rate of Take (Proportion of Total Take) of Incidentally Caught Fish in Research Gill Nets Set to Catch Herring.....	4-16

